

Ultrafast all-optical control in photonic crystal cavities using nonlinear absorption

C.-S. Kee*, Y. L. Lee, D.-K. Ko, and J. Lee

Advanced Photonics Research Institute, Gwangju Institute of Science and Technology, Gwangju 500-712, Republic of Korea

H.-G. Choi, J.-E. Kim, and H. Y. Park

Department of Physics, Korea Advanced Institute of Science and Technology, Daejeon 305-701, Republic of Korea

Optical nonlinearities of semiconductors can offer interesting effects on the properties of photonic crystals based on semiconductors [1]. In particular, the nonlinear absorption such as two-photon absorption can give an opportunity to control optically the properties of semiconductor photonic crystals. Moreover, a fast response time of free carriers generated by two-photon absorption can offer rapid control in semiconductor photonic crystals. Thus the nonlinear absorption can be useful for achieving ultrafast all-optical control in photonic crystal devices. In this presentation, we theoretically show that the properties of AlGaAs photonic crystal cavity can be optically controlled by two-photon absorption. The measured free carrier induced refractive index change as a function of pump power [2] was employed in the calculations. The resonant wavelength λ_{re} exhibits a blueshift when the pump power increases. The modulation depth, defined as $|(T_{\text{pump-off}} - T_{\text{pump-on}}) / T_{\text{pump-off}}|$, can reach about 90 % when the pump power is 2.5 mJ/cm^2 . The modulation time is predicted to be faster than 10 ps.




Figure 1. Transmission spectra for TM mode whose magnetic field is parallel to the axis of a hole through AlGaAs photonic crystal cavity (a) and spatial distributions of magnetic field for $\lambda_{re, \text{pump-off}}$ when the pump laser with power of 2.5 mJ/cm^2 is off and on (b). Schematic top view of AlGaAs photonic crystal cavity and its structural parameters are shown in (c).

Reference

*Email: cskee@gist.ac.kr

[1] A. Hache and M. Bourgeois, *Applied Physics Letters*, **77**, 4089 (2000).

[2] A. D. Bristow, D. O. Kundys, A. Z. Garcia-Deniz, J. P. R. Wells, A. M. Fox, M. S. Skolnick, D. M. Whittaker, A. Tahraoui, T. F. Krauss, and J. S. Roberts, *Journal of Applied Physics*, **96**, 4729 (2004).